

To view this information online or in other languages, go to www.meducation.com.

meducation®



0 1 U 2 4 3 G N



**UNIVERSITY
MEDICAL CENTER**

University Medical Center
123 Main Street
Anytown, NC 12345
212-555-2121





Biaxin Oral Suspension 125 mg/5 mL


This medicine is used to treat infection.

How to take medicine

Take the medicine by mouth twice a day with meals - breakfast and dinner.

Drink 5 mL each time.

			
Breakfast	Lunch	Dinner	Bedtime
5 mL		5 mL	



Use the medicine for a total of 7 days.

Instructions

Take the medicine with food.

Keep this medicine at room temperature.

Shake the bottle well before using the medicine.

Avoid drinking grapefruit juice while taking this medicine.

After you have finished taking your medicine for the number of days indicated, please throw away any remaining unused medicine.

It is important that you keep taking each dose of this medicine on time even if you are feeling well.

If you forget to take a dose on time, take it as soon as you remember. If it is almost time for the next dose, do not take the missed dose. Return to your normal dosing schedule. Do not take 2 doses of this medicine at one time.

Please tell your doctor or pharmacist if you take other prescription or over-the-counter medicines, vitamins, herbal medicines, or anything else for your health.



Cautions

If you have ever had a severe allergic reaction to another medicine, please check with your doctor or pharmacist to determine if you should avoid taking this medicine. Symptoms of a severe allergic reaction include difficulty breathing, skin rash, itching, swelling, or severe dizziness.

Do not use the medication any more than instructed.

Do not drink beverages containing alcohol while on this medicine.

Please tell your doctor if you have moderate to severe diarrhea while on this medicine. Do not treat the diarrhea with over-the-counter diarrhea medicine.

Tell the doctor or pharmacist if you are pregnant, planning to be pregnant, or breastfeeding.

Check with your doctor or pharmacist about potential interactions with your other medications.

Do not start or stop any other medicines without first speaking to your doctor or pharmacist.

Do not share this medicine with anyone who has not been prescribed this medicine.

Always keep medicine out of reach of children and pets.

Side Effects

- diarrhea
- headaches
- liver problems
- nausea
- skin irritation such as redness, itching, rash, or burning
- stomach upset or abdominal pain
- changes in taste or unpleasant taste
- vomiting

These are some potential side effects from this medicine. Please speak with your doctor about what you should do if you experience these or other side effects from your medication.

An allergic reaction to this medicine is uncommon. But if you experience difficulty breathing, skin rash, itching, swelling, or severe dizziness, seek medical care immediately.

Refill

You do not have any refills for this medicine.

Please speak with your doctor, nurse, or pharmacist if you have any questions about this medicine.

IMPORTANT NOTE: This information is being provided to you as a supplement to other written instructions and package inserts provided by your doctor or pharmacist. This document contains a summary of your medication instructions and may not include all the product warnings or other important information about your medication. Please speak with your doctor or pharmacist about any question you may have about your medication or any unexpected symptoms. Always follow their advice and instructions.



To view this information online or in other languages, go to www.meducation.com.

meducation®



University Medical Center
123 Main Street
Anytown, NC 12345
212-555-2121





Foradil Aerolizer 12 mcg/inh

This medicine is used to treat chronic lung disease.

How to take medicine

Use the medicine twice a day.

Inhale one (1) puff each time.

			
Morning	Noon	Evening	Bedtime
1 puff		1 puff	



You should keep taking this medicine until you are told to stop.

Instructions

The capsule needs to be placed into your inhaler before each use. Please follow the instructions for your inhaler carefully. Do not swallow the capsule.

Keep this medicine at room temperature. Protect from moisture and high humidity.

Ask your doctor, nurse or pharmacist to show you how to use this medicine correctly.

The medicine is a very fine powder. Some patients may not feel or taste the medicine. Do not take an extra dose of medicine if you do not taste or feel the medicine being delivered.

It is important that you keep taking each dose of this medicine on time even if you are feeling well.

If you forget to take a dose on time, take it as soon as you remember. If it is almost time for the next dose, do not take the missed dose. Return to your normal dosing schedule. Do not take 2 doses of this medicine at one time.

Please tell your doctor or pharmacist if you take other prescription or over-the-counter medicines, vitamins, herbal medicines, or anything else for your health.

Do not share this medicine with anyone who has not been prescribed this medicine.

Cautions

Do not take the medicine more than twice during 24 hours.

Do not use the medication any more than instructed.

If the medicine causes increased wheezing or difficulty breathing, stop the medicine immediately and seek immediate medical care.

Your ability to stay alert or to react quickly may be impaired by this medicine. Do not drive or operate machinery until you know how this medicine will affect you.

Tell the doctor or pharmacist if you are pregnant, planning to be pregnant, or breastfeeding.

Do not use this medicine during an asthma attack.

This medicine will not stop an asthma attack once it has started.

Do not start or stop any other medicines without first speaking to your doctor or pharmacist.

If you have difficulty using your medicine, please tell your doctor, nurse or pharmacist.

Some patients taking this medicine have experienced serious side effects. Additional information approved by the U.S. Food & Drug Administration (FDA) is available from your pharmacist for this medicine. Please review the document with your pharmacist to understand the risks associated with this medicine to prevent potential serious side effects.

Always keep medicine out of reach of children and pets.

Side Effects

- agitated feeling or trouble sleeping
- rarely, a severe episode of asthma may occur
- dizziness
- dry mouth
- lack of energy and tiredness
- increased risk for an infection
- nausea
- rapid heartbeat
- shakiness
- stomach upset or abdominal pain
- changes in taste or unpleasant taste

These are some potential side effects from this medicine. Please speak with your doctor about what you should do if you experience these or other side effects from your medication.

An allergic reaction to this medicine is uncommon. But if you experience difficulty breathing, skin rash, itching, swelling, or severe dizziness, seek medical care immediately.

Refill

You may refill this medicine twice (2).

Please speak with your doctor, nurse, or pharmacist if you have any questions about this medicine.

IMPORTANT NOTE: This information is being provided to you as a supplement to other written instructions and package inserts provided by your doctor or pharmacist. This document contains a summary of your medication instructions and may not include all the product warnings or other important information about your medication. Please speak with your doctor or pharmacist about any question you may have about your medication or any unexpected symptoms. Always follow their advice and instructions.



To view this information online or in other languages, go to www.meducation.com.

meducation®



0 1 U 2 4 3 G N



**UNIVERSITY
MEDICAL CENTER**

University Medical Center
123 Main Street
Anytown, NC 12345
212-555-2121



Prednisone Tablet 5 mg





This medicine is used to treat asthma.

How to take medicine

Your dosage schedule for this medicine is shown below.

Please take the number of pills shown for each day.

To help you keep track of the medicine you have taken, please cross out the box as you take each dose.

	 Breakfast	 Lunch	 Dinner	 Bedtime
Day 1	5	-	-	-
Day 2	5	-	-	-
Day 3	4	-	-	-
Day 4	4	-	-	-
Day 5	3	-	-	-
Day 6	3	-	-	-
Day 7	2	-	-	-
Day 8	2	-	-	-
Day 9	1	-	-	-
Day 10	1	-	-	-

Stop taking the medicine after the last day of your dosage schedule.

Use the medicine for a total of 10 days.

Instructions

Swallow the medicine without crushing or chewing it.

Take the medicine with food.

Keep the medicine at room temperature. Avoid heat and direct light.

It is important that you keep taking each dose of this medicine on time even if you are feeling well.

If you forget to take a dose on time, take it as soon as you remember. If it is almost time for the next dose, do not take the missed dose. Return to your normal dosing schedule. Do not take 2 doses of this medicine at one time.

Please tell your doctor or pharmacist if you take other prescription or over-the-counter medicines, vitamins, herbal medicines, or anything else for your health.

Do not suddenly stop taking this medicine without checking with your doctor.

This medicine may affect your blood sugar levels. If you have diabetes, tell your doctor about any changes in your blood sugar levels before adjusting your diabetes medicine.

This medicine may affect the strength of your bones. If you have or are at increased risk for developing osteoporosis (weakening of the bones), your doctor may recommend adding foods containing calcium and vitamin D while on this medicine. Please talk to your doctor for more information.

Cautions

If you have ever had a severe allergic reaction to another medicine, please check with your doctor or pharmacist to determine if you should avoid taking this medicine. Symptoms of a severe allergic reaction include difficulty breathing, skin rash, itching, swelling, or severe dizziness.

Do not use the medication any more than instructed.

This medicine may reduce your body's ability to fight infections. Try to avoid contact with people with colds, flu or other infections.

Tell the doctor or pharmacist if you are pregnant, planning to be pregnant, or breastfeeding.

Check with your doctor or pharmacist about potential interactions with your other medications.

Please tell any doctor or dentist that you are on this medicine before they provide any medical treatment, dental care, or surgery.

Do not start or stop any other medicines without first speaking to your doctor or pharmacist.

Do not share this medicine with anyone who has not been prescribed this medicine.

Always keep medicine out of reach of children and pets.

Side Effects

- agitated feeling or trouble sleeping
- increased risk of bleeding
- changes in memory, mood, or thinking
- depression or feeling sad
- swelling of the legs, feet, and hands
- headaches
- high blood sugar
- high blood pressure
- muscle aches, spasms or abnormal movements
- nausea
- rapid heartbeat
- skin irritation such as redness, itching, rash, or burning
- shakiness
- stomach upset or abdominal pain
- vomiting
- weight gain

These are some potential side effects from this medicine. Please speak with your doctor about what you should do if you experience these or other side effects from your medication.

An allergic reaction to this medicine is uncommon. But if you experience difficulty breathing, skin rash, itching, swelling, or severe dizziness, seek medical care immediately.

Refill

You do not have any refills for this medicine.

Please speak with your doctor, nurse, or pharmacist if you have any questions about this medicine.

IMPORTANT NOTE: This information is being provided to you as a supplement to other written instructions and package inserts provided by your doctor or pharmacist. This document contains a summary of your medication instructions and may not include all the product warnings or other important information about your medication. Please speak with your doctor or pharmacist about any question you may have about your medication or any unexpected symptoms. Always follow their advice and instructions.

To view this information online or in other languages, go to www.meducation.com.

meducation®



0 1 U 2 4 3 G N



**UNIVERSITY
MEDICAL CENTER**

University Medical Center
123 Main Street
Anytown, NC 12345
212-555-2121





Lumigan Ophthalmic Solution 0.03%

This medicine is used to treat glaucoma.

How to take medicine

Use the medicine once a day.

Use one (1) drop each time.

			
Morning	Noon	Evening	Bedtime
1 drop			



Put the medicine into both eyes.

Use the medicine everyday until your doctor tells you to stop.

Instructions

If you use contact lenses, remove the contacts before using the medicine and then wait 15 minutes after using the medicine before inserting them.

Keep the medicine at room temperature. Avoid heat and direct light.

If you forget to take a dose on time, take it as soon as you remember. If it is almost time for the next dose, do not take the missed dose. Return to your normal dosing schedule. Do not take 2 doses of this medicine at one time.

Please tell your doctor or pharmacist if you take other prescription or over-the-counter medicines, vitamins, herbal medicines, or anything else for your health.

If you are using another eye medicine, wait at least 5 minutes before using the other medicine.

Cautions

If you have ever had a severe allergic reaction to another medicine, please check with your doctor or pharmacist to determine if you should avoid taking this medicine. Symptoms of a severe allergic reaction include difficulty breathing, skin rash, itching, swelling, or severe dizziness.

Do not use the medication any more than instructed.

This medicine may cause blurry vision.

Tell the doctor or pharmacist if you are pregnant, planning to be pregnant, or breastfeeding.

Check with your doctor or pharmacist about potential interactions with your other medications.

Do not share this medicine with anyone who has not been prescribed this medicine.

Always keep medicine out of reach of children and pets.

Side Effects

- blurry vision
- discomfort in the eyes
- eye lashes growing longer
- irritation of the eyes
- color changes in the eyes and eyelids
- redness of eyes
- blurring or changes of vision

These are some potential side effects from this medicine. Please speak with your doctor about what you should do if you experience these or other side effects from your medication.

An allergic reaction to this medicine is uncommon. But if you experience difficulty breathing, skin rash, itching, swelling, or severe dizziness, seek medical care immediately.

Refill

You may refill this medicine twice (2).

Please speak with your doctor, nurse, or pharmacist if you have any questions about this medicine.

IMPORTANT NOTE: This information is being provided to you as a supplement to other written instructions and package inserts provided by your doctor or pharmacist. This document contains a summary of your medication instructions and may not include all the product warnings or other important information about your medication. Please speak with your doctor or pharmacist about any question you may have about your medication or any unexpected symptoms. Always follow their advice and instructions.

